

## EDITORIAL

# Arteriosclerosis

## A Primer for “In Focus” Reviews on Arterial Stiffness

Gary F. Mitchell, Janet T. Powell

**A**rteriosclerosis, a descriptive term for thickening and hardening of the arterial wall, accompanies aging and is associated with major adverse impact on the cardiovascular system and a number of other diseases that are now known to have a strong underlying microvascular component to their pathogenesis, including chronic kidney disease, vascular dementia, and Alzheimer disease.<sup>1,2</sup> Although a multitude of studies have attempted to identify genetic contributions to arterial stiffening in humans, the results have been largely inconclusive, in part due to lack of appreciation of both the population epidemiology of arterial stiffness and the variety of measurement methods and arterial beds used to assess arterial stiffness.<sup>3</sup> Recently, 2 separate groups have capitalized on the UK Biobank study to report genome-wide association studies of finger plethysmography measurements of arterial stiffness index, with emerging indications that collagen gene *COL4A2* variants may be associated with arterial stiffness index.<sup>4,5</sup> However, finger plethysmography does not provide a robust measure of the stiffness of the central elastic arteries, which is of critical importance to cardiac function. Progress in the pathobiology of arterial stiffness depends on designing experimental studies that consider the challenges posed by the epidemiology and measurement of arterial stiffness. Therefore, the purpose of the forthcoming series of “In Focus” reviews is to identify these challenges for vascular biologists together with their clinical and genetic colleagues.

Arterial stiffness or arteriosclerosis is often associated with but is distinct from atherosclerosis, which is a patchy intimal abnormality that likely represents a consequence rather than a cause of arteriosclerosis.<sup>6,7</sup> Arteriosclerosis arises as a result of increased production or progressively greater engagement of stiffer load-bearing elements in the arterial wall, such as collagen.

Stiffening of the aorta increases pulsatile load on the left ventricle, leading to hypertrophy and diastolic dysfunction, which can progress to heart failure, often with preserved left ventricular ejection fraction.<sup>8–10</sup> Aortic stiffening reduces the normal impedance gradient between aorta and branching conduits and thereby increases pulsatile pressure and flow in downstream conduits, such as the carotid and renal arteries, leading to transmission of excessive pulsatility into the fragile microcirculation, particularly in high-flow organs such as the brain and kidneys.<sup>11–13</sup> In these low impedance vascular beds, low precapillary resistance allows excessive pulsatile energy that enters the local conduit artery to be transmitted directly into the microcirculation, where it causes microvascular damage, remodeling and repair, leading to target organ damage and dysfunction.

Risk factors for arterial stiffening other than age and distending pressure remain relatively obscure.<sup>6</sup> Though once thought to represent a consequence of long-standing hypertension, it is now clear that aortic stiffening precedes and contributes to the pathogenesis of hypertension.<sup>14</sup> Arteriosclerosis arises in part as a consequence of a lifelong progressive increase in aortic diameter, which accommodates an increase in cardiac output associated with normal somatic growth in early life and progressive pathological obesity from adulthood onward.<sup>15</sup> While aortic remodeling maintains the balance between flow and diameter, it may accelerate deterioration of elastic elements in the aortic wall. The complement of medial elastic fibers in the aortic wall is determined by a complex interplay of genetic and environmental factors and is completed early in life at the equivalent of the toddler stage in various animal models and humans. The complex gene program that is required to produce

**Key Words:** Editorials ■ arteriosclerosis ■ epidemiology ■ hypertension ■ plethysmography ■ vascular dementia

Correspondence to: Gary F. Mitchell, MD, Cardiovascular Engineering, Inc, 1 Edgewater Dr, Suite 201, Norwood, MA 02062. Email garyfmitchell22@gmail.com  
For Sources of Funding and Disclosures, see page 1026.

© 2020 American Heart Association, Inc.

*Arterioscler Thromb Vasc Biol* is available at [www.ahajournals.org/journal/atvb](http://www.ahajournals.org/journal/atvb)

medial elastic fibers is subsequently silenced,<sup>16,17</sup> resulting in a fixed complement of elastic fibers that must last a lifetime.<sup>18</sup>

Progressive remodeling to a larger diameter of this fixed pool of elastic fibers in the aortic media leads to thinning and increased elastic fiber stress and transfers progressively greater load bearing to much stiffer collagen fibers. As the wall stiffens, pulse wave velocity increases. Remodeling to a larger diameter can limit the increase in pulse pressure (the difference between systolic and diastolic pressure) associated with a given increase in wall stiffness because characteristic impedance of the aorta ( $Z_c$ ), which defines the change in pressure ( $dP$ ) produced by a given change in flow ( $dQ$ ) in the aorta ( $Z_c=dP/dQ$ ), has a strong inverse association with lumen diameter. However, progressive stiffening eventually overwhelms diameter remodeling at around midlife, leading to a progressive increase in systolic and pulse pressure thereafter, particularly in women.<sup>15,19</sup> The increase in pulse pressure plays a major role in the pathogenesis of isolated or predominant systolic hypertension, which is highly prevalent after midlife and is often difficult to control adequately.<sup>20</sup>

The combination of high risk associated with aortic arteriosclerosis plus the epidemic prevalence of increased stiffness in middle-aged and older people implies that the burden of disease attributable to stiffness will rise considerably over the next 2 decades as the population ages if we do not find ways to prevent or reverse excessive stiffness.<sup>21,22</sup> While age-related stiffening is highly prevalent, it is not inevitable, underscoring the need for additional research on pathogenesis and vascular biology of stiffening and factors that promote healthy vascular aging.<sup>23</sup>

In this context, *ATVB* has commissioned a series of brief reviews that aims to underscore the critical importance of aortic arteriosclerosis in the pathogenesis of cardiovascular disease while outlining what we know and what we need to better understand about pathogenesis and consequences of arteriosclerosis. Wilkinson et al<sup>7</sup> present arterial stiffness in the clinical context, while Yu and McEniery present a brief review of the epidemiology of stiffness. Schutte et al provide an overview of ethnic differences in aortic stiffness. Segers et al describe various techniques that have been developed to assess arteriosclerosis in humans while Butlin et al address the unique challenges of performing stiffness measurements in small animals. Laurent et al summarize current understanding of various factors, including smooth muscle cell biology, that contribute to stiffening of the arterial wall while Chen et al look specifically at factors associated with arterial calcification.

We trust that this series of reviews will stimulate high-quality aspirational research that aims to elucidate

factors that contribute to arterial stiffening, its consequences and reversal.

## ARTICLE INFORMATION

### Affiliations

From Cardiovascular Engineering, Inc, Norwood, MA (G.F.M.); and Faculty of Medicine, Department of Surgery & Cancer, Imperial College London, United Kingdom (J.T.P.).

### Acknowledgments

Drs Mitchell and Powell conceived the article and edited the article for intellectual content.

### Sources of Funding

Dr Mitchell was supported by grants from the National Institutes for Health, National Heart Lung and Blood Institute.

### Disclosures

Dr Mitchell is the owner of Cardiovascular Engineering, Inc, a small business that designs and develops devices that are used to measure arterial stiffness and has received consulting fees or grants from Novartis, Merck, Bayer, and Servier. Dr Powell is an associate editor of *Arteriosclerosis Thrombosis and Vascular Biology*.

## REFERENCES

- Dagenais GR, Leong DP, Rangarajan S, Lanas F, Lopez-Jaramillo P, Gupta R, Diaz R, Avezum A, Wielgosz A, Parambath SR, et al. Variations in common diseases, hospital admissions, and deaths in middle-aged adults in 21 countries from five continents (PURE): a prospective cohort study. *Lancet*. 2020;395:785–794. doi: 10.1016/S0140-6736(19)32007-0
- Mitchell GF. Aortic stiffness, pressure and flow pulsatility, and target organ damage. *J Appl Physiol* (1985). 2018;125:1871–1880. doi: 10.1152/jappphysiol.00108.2018
- Logan JG, Engler MB, Kim H. Genetic determinants of arterial stiffness. *J Cardiovasc Transl Res*. 2015;8:23–43. doi: 10.1007/s12265-014-9597-x
- Fung K, Ramirez J, Warren HR, Aung N, Lee AM, Tzani E, Petersen SE, Munroe PB. Genome-wide association study identifies loci for arterial stiffness index in 127,121 UK Biobank participants. *Sci Rep*. 2019;9:9143. doi: 10.1038/s41598-019-45703-0
- Zekavat SM, Aragam K, Emdin C, Khara AV, Klarin D, Zhao H, Natarajan P. Genetic association of finger photoplethysmography-derived arterial stiffness index with blood pressure and coronary artery disease. *Arterioscler Thromb Vasc Biol*. 2019;39:1253–1261. doi: 10.1161/ATVBAHA.119.312626
- Cecelja M, Chowienczyk P. Dissociation of aortic pulse wave velocity with risk factors for cardiovascular disease other than hypertension: a systematic review. *Hypertension*. 2009;54:1328–1336. doi: 10.1161/HYPERTENSIONAHA.109.137653
- Wilkinson IB, McEniery CM, Cockcroft JR. Arteriosclerosis and atherosclerosis: guilty by association. *Hypertension*. 2009;54:1213–1215. doi: 10.1161/HYPERTENSIONAHA.109.142612
- Chirinos JA, Khan A, Bansal N, Dries DL, Feldman HI, Ford V, Anderson AH, Kallem R, Lash JP, Ojo A, et al; CRIC Study Investigators; CRIC Study Investigators. Arterial stiffness, central pressures, and incident hospitalized heart failure in the chronic renal insufficiency cohort study. *Circ Heart Fail*. 2014;7:709–716. doi: 10.1161/CIRCHEARTFAILURE.113.001041
- Ohyama Y, Ambale-Venkatesh B, Noda C, Kim JY, Tanami Y, Teixeira-Tura G, Chugh AR, Redheuil A, Liu CY, Wu CO, et al. Aortic arch pulse wave velocity assessed by magnetic resonance imaging as a predictor of incident cardiovascular events: the MESA (Multi-Ethnic Study of Atherosclerosis). *Hypertension*. 2017;70:524–530. doi: 10.1161/HYPERTENSIONAHA.116.08749
- Tsao CW, Lyass A, Larson MG, Levy D, Hamburg NM, Vita JA, Benjamin EJ, Mitchell GF, Vasani RS. Relation of central arterial stiffness to incident heart failure in the community. *J Am Heart Assoc*. 2015;4:e002189. doi: 10.1161/JAHA.115.002189
- Cooper LL, Woodard T, Sigurdsson S, van Buchem MA, Torjesen AA, Inker LA, Aspelund T, Eiriksdottir G, Harris TB, Gudnason V, et al. Cerebrovascular damage mediates relations between aortic stiffness and memory. *Hypertension*. 2016;67:176–182. doi: 10.1161/HYPERTENSIONAHA.115.06398
- Mitchell GF, van Buchem MA, Sigurdsson S, Gotal JD, Jonsdottir MK, Kjartansson Ó, Garcia M, Aspelund T, Harris TB, Gudnason V, et al. Arterial

- stiffness, pressure and flow pulsatility and brain structure and function: the Age, Gene/Environment Susceptibility-Reykjavik study. *Brain*. 2011;134(pt 11):3398–3407. doi: 10.1093/brain/awr253
13. Woodard T, Sigurdsson S, Gotlib JD, Torjesen AA, Inker LA, Aspelund T, Eiriksdottir G, Gudnason V, Harris TB, Launer LJ, et al. Mediation analysis of aortic stiffness and renal microvascular function. *J Am Soc Nephrol*. 2015;26:1181–1187. doi: 10.1681/ASN.2014050450
  14. Kaess BM, Rong J, Larson MG, Hamburg NM, Vita JA, Levy D, Benjamin EJ, Vasani RS, Mitchell GF. Aortic stiffness, blood pressure progression, and incident hypertension. *JAMA*. 2012;308:875–881. doi: 10.1001/2012.jama.10503
  15. Mitchell GF, Wang N, Palmisano JN, Larson MG, Hamburg NM, Vita JA, Levy D, Benjamin EJ, Vasani RS. Hemodynamic correlates of blood pressure across the adult age spectrum: noninvasive evaluation in the Framingham Heart Study. *Circulation*. 2010;122:1379–1386. doi: 10.1161/CIRCULATIONAHA.109.914507
  16. Johnson DJ, Robson P, Hew Y, Keeley FW. Decreased elastin synthesis in normal development and in long-term aortic organ and cell cultures is related to rapid and selective destabilization of mRNA for elastin. *Circ Res*. 1995;77:1107–1113. doi: 10.1161/01.res.77.6.1107
  17. Ott CE, Grünhagen J, Jäger M, Horbelt D, Schwill S, Kallenbach K, Guo G, Manke T, Knaus P, Mundlos S, et al. MicroRNAs differentially expressed in postnatal aortic development downregulate elastin via 3' UTR and coding-sequence binding sites. *PLoS One*. 2011;6:e16250. doi: 10.1371/journal.pone.0016250
  18. Powell JT, Vine N, Crossman M. On the accumulation of D-aspartate in elastin and other proteins of the ageing aorta. *Atherosclerosis*. 1992;97:201–208. doi: 10.1016/0021-9150(92)90132-z
  19. Mitchell GF, Gudnason V, Launer LJ, Aspelund T, Harris TB. Hemodynamics of increased pulse pressure in older women in the community-based Age, Gene/Environment Susceptibility-Reykjavik Study. *Hypertension*. 2008;51:1123–1128. doi: 10.1161/HYPERTENSIONAHA.107.108175
  20. Franklin SS, Jacobs MJ, Wong ND, L'Italien GJ, Lapuerta P. Predominance of isolated systolic hypertension among middle-aged and elderly US hypertensives: analysis based on National Health and Nutrition Examination Survey (NHANES) III. *Hypertension*. 2001;37:869–874. doi: 10.1161/01.hyp.37.3.869
  21. Mitchell GF, Guo CY, Benjamin EJ, Larson MG, Keyes MJ, Vita JA, Vasani RS, Levy D. Cross-sectional correlates of increased aortic stiffness in the community: the Framingham Heart Study. *Circulation*. 2007;115:2628–2636. doi: 10.1161/CIRCULATIONAHA.106.667733
  22. Mitchell GF, Hwang SJ, Vasani RS, Larson MG, Pencina MJ, Hamburg NM, Vita JA, Levy D, Benjamin EJ. Arterial stiffness and cardiovascular events: the Framingham Heart Study. *Circulation*. 2010;121:505–511. doi: 10.1161/CIRCULATIONAHA.109.886655
  23. Niiranen TJ, Lyass A, Larson MG, Hamburg NM, Benjamin EJ, Mitchell GF, Vasani RS. Prevalence, correlates, and prognosis of healthy vascular aging in a western community-dwelling cohort: the Framingham Heart Study. *Hypertension*. 2017;70:267–274. doi: 10.1161/HYPERTENSIONAHA.117.09026